

# Cal Poly Department of Mathematics

## Puzzle of the Week

Feb 13 - 19, 2009

Suggested by Kent Morrison:

Fix three points in the plane at  $(1, 0)$ ,  $(-1, 0)$ ,  $(0, h)$  where  $h > 0$ . Find the point(s)  $(x, y)$  the sum of whose distances to these three points is minimized.

*Solutions should be submitted to Morgan Sherman:*

*Dept. of Mathematics, Cal Poly*

*Email: sherman1 -AT- calpoly.edu*

*Office: bldg 25 room 310*

*before next Friday. Those with correct and complete solutions will have their names listed in next week's email announcement. Anybody is welcome to make a submission.*

*Solution:* The point is given by

$$P = (0, t) \text{ where } t = \min \left\{ h, \frac{1}{\sqrt{3}} \right\}.$$

Note that for  $h > 3^{-\frac{1}{2}}$  this point is independent of  $h$ !

Most solutions begin “clearly  $P$  lies on the (positive)  $y$ -axis...”. I can't really argue with that, but I will include a geometric argument to justify this anyway. (Jeff Liese gave a different argument, using Lagrange multipliers, which proves that of all triangles with fixed base and height the isosceles minimizes the perimeter).

Let  $F_1 = (-1, 0)$ ,  $F_2 = (1, 0)$ ,  $H = (0, h)$  and let  $P$  be a point which minimizes the sum of the distances to these points. Consider the (unique) ellipse passing through  $P$  and having  $F_1, F_2$  as foci. By its definition every point on this ellipse has the same sum of distances to  $F_1, F_2$ . Now if  $H$  lies inside this ellipse, then by comparing with the smaller ellipse passing through  $H$  we see that replacing  $P$  with  $H$  gives a smaller sum of distances to  $F_1, F_2, H$ , a contradiction. Otherwise we note that the point on the ellipse intersecting the positive  $y$  axis is the closest to  $H$  and therefore  $P$  must be this point.

Now write  $P = (0, t)$  with  $t \in [0, h]$  and set

$$g(t) = h - t + 2\sqrt{1 + t^2}, \quad 0 \leq t \leq h.$$

We consider the critical points of  $g$ :

$$g'(t) = -1 + \frac{2t}{\sqrt{1 + t^2}} = 0 \iff t = \pm \frac{1}{\sqrt{3}}$$

So if  $h \geq 1/\sqrt{3}$  then we have one critical point on  $[0, h]$ , namely  $t = 1/\sqrt{3}$ , and since  $g''(1/\sqrt{3}) > 0$  this is the unique minimum.

On the other hand if  $0 < h < 1/\sqrt{3}$  then there are no critical points so we must test the end points:

$$g(0) = h + 2, \quad g(h) = 2\sqrt{1 + h^2}.$$

By squaring both sides and doing a little algebra we find  $g(h) \leq g(0)$  when  $h < 1/\sqrt{3}$ . Hence our final answer.