

# Cal Poly Department of Mathematics

## Puzzle of the Week

May 22 - 28, 2009

In the Cleveland Cavaliers' first game of the 2008-09 season we witnessed LeBron James make 4 out of his 8 free throw attempts, thus ending the game shooting 50% from the stripe. After the last game of the season LeBron finished with a season average of 78% of free thows made.

Question: In going from 50 to 78 percent must LeBron have averaged exactly 75% at some point in time after game 1 and by the end of the season?

*Solutions should be submitted to Morgan Sherman:*

*Dept. of Mathematics, Cal Poly  
Email: sherman1 -AT- calpoly.edu  
Office: bldg 25 room 310*

*before next Friday. Those with correct and complete solutions will have their names listed in next week's email announcement. Anybody is welcome to make a submission.*

*Solution:* This problem comes from the 2004 Putnam exam.

The answer is yes, LeBron did have to actually hit exactly 75% of his free throws at some point. To see this suppose that he managed to go from under 75% to over 75% without hitting exactly 75%. Consider the free throw which sent him over the threshold. If  $a$  denotes the number of made free throws just before this point, and  $b$  denotes total number of attempts at that time, then those numbers satisfy

$$\frac{a}{b} < \frac{3}{4} < \frac{a+1}{b+1}$$

But then  $4a < 3b < 4a + 1$  which is impossible since  $a$  and  $b$  are integers.

Note that if we replace  $3/4$  with  $(k-1)/k$  then the same reasoning will still hold. So LeBron would also have made exactly  $2/3$  of his shots at some point. And any basketball player who goes from 0% to 91% (say) will have made exactly  $1/2, 2/3, 3/4, 4/5, 5/6, 6/7, 7/8, 8/9, 9/10$  and  $10/11$  of his or her shots at some point.

Thanks for following the Cal Poly "Puzzle of the Week" this year. It will return in the Fall, so stay tuned..